

Prevalence of Work Related Musculoskeletal Disorders in Farmers of Gujarat

Dr. Bhavana Gadhavi¹, Dr. Yagna Shukla²

¹PhD Scholar, Gujarat University, Ahmadabad, Gujarat.

²Senior Lecturer at Government Physiotherapy Hospital, Civil Hospital Campus, Girdharnagar, Ahmedabad-380016.

Corresponding Author: Dr. Bhavana Gadhavi

ABSTRACT

Introduction: Musculoskeletal Disorders (MSDs) are a global problem and have a comprehensive impact on health and economy of a country. Work-related musculoskeletal disorders (WMSDs) are accumulative disorders that are most frequently found in agricultural farmers. Farming is a physically arduous occupation. Presence of musculoskeletal disorders among these farmers will result in reduction in their work capacity which may reflect in reduction in economic contribution from farming sector. The present study was conducted to find out the prevalence of musculoskeletal disorders among farmers.

Methodology: All adult farmers between the ages of 18 to 70 years were included. 947 workers were selected randomly and the prevalence of musculoskeletal disorders was determined by self developed questionnaire. Result The most common MSDs were found to be in the Lower back (61%) followed by Knee (33%), Shoulder (31%), Neck(29%), Upper back (28%), Elbow (15%) , Hip (13%), Wrist and hand (12%), and Ankle and Foot (11%).

Conclusion: The prevalence of MSDs in farmers is high in farmers of Gujarat with highest prevalence in Lower Back.

Key words: Work-related musculoskeletal disorders, Farmers, Gujarat

INTRODUCTION

Musculoskeletal Disorders (MSDs) are a global problem and have a comprehensive impact on health and economy of a country. [1] Musculoskeletal disorders are defined as a “Group of disorders that affect the musculoskeletal system including the nerves, muscles, tendons and joints and supporting structures such as inter-vertebral discs etc.” [4] Work-related musculoskeletal disorders (WMSDs) are accumulative disorders that are most frequently found in agricultural farmers. [2] Farming is a physically arduous occupation and this places farm workers at potential risk of musculoskeletal disorders such as osteoarthritis (OA) of the hip and knee, low

back pain (LBP), neck and upper limb complaints, and hand–arm vibration syndrome (HAVS). [3] It may result in pain, damage to musculoskeletal structures, poor health as well as poor quality of life and reduced productivity. [5] They are the most common cause of severe persisting pain and disability, and are currently reported to be affecting hundreds of millions of people across the globe. [6-8] It often occurs in an individual, when the work load exceeds the capacity that his / her locomotor apparatus can bear. WMSDs may occur as a result of acute injuries from one-time trauma or multiple traumas such as repetitive motion, excessive force, sustained abnormal postures, prolonged squatting and standing

in the course of work. [15] The disorder takes on a more serious dimension when it becomes chronic; nearly 25 percent of the affected adults are identified as having chronic Musculoskeletal (MS) impairment, which is equally prevalent in both developed as well as developing countries. [10,11]

Ergonomics is the scientific discipline concerned with the understanding of the interactions among humans and other elements of a system and the profession that applies theory, principles, data, and methods to design in order to optimize human well-being and overall system performance. In the design of work and everyday life situations, the focus of ergonomics is a man. Unsafe, unhealthy, uncomfortable or inefficient situations at work or in everyday life are avoided by taking account of the physical and psychological capabilities and limitations of a human. [9]

The major work related risk factors associated with LBP have been identified as poor/awkward work postures, bending, lifting and physically strenuous work. [12-14] Based on the International Labour Organization, globally 74% of the agricultural workers live in Asia and Pacific regions. [16] In India, as 68% of the populations live in rural areas, agriculture based activities play a major role in improving the rural economy of the country. [17,18] Agriculture and its related activities have provided nearly 60% of the employment opportunities in India. [19] WMSDs are found to be associated with absenteeism, loss of productivity and economic loss to the worker, industry and the nation at large. [20] There are several studies globally showing that the agriculture workers have a higher risk of developing MSDs than any other group of people.

However, there are scarcities of studies that focus on the country's farming community, which constitutes more than 58 percent of the Indian work force. [21] Presence of musculoskeletal disorders among these farmers will result in reduction in their work capacity which may reflect in

reduction in economic contribution from farming sector. After a thorough literature search, it had been found that the prevalence of musculoskeletal disorders in various regions of body among farming community in this part of the country is not well documented. Hence, the present study was conducted to find out the prevalence of musculoskeletal disorders among farmers.

MATERIALS AND METHODS

Ethical approval was obtained from Institutional Ethics Committee. A community based cross-sectional study was conducted from December 2017 to September 2019.

All adult farmers between the ages of 18 to 70 years were included. Part-time farmers (i.e. people who were also doing some other jobs besides farming e.g. rickshaw pullers, carpenters, other labour workers); subjects with diagnosed congenital skeletal deformities or deformities due to fractures; subjects with any diagnosed psychiatric illness; who were known to have spinal fracture resulting from tumours, infections or any major trauma to the spine or having diagnosed neurological problems were excluded. Unwilling subjects were also excluded. Informed written consent was taken from all included farmers.

947 workers were selected randomly and the prevalence of musculoskeletal disorders was determined by self developed questionnaire. In this questionnaire the musculoskeletal system is divided to 9 regions such as neck, shoulders, hip, upper back, knees, wrists/hands, ankles/feet, low back and knee. The researcher completed the questionnaires by interview with workers. In very explicit and simple terms respondents were asked if they had experienced any musculoskeletal discomfort in any of the joints in their body which prevented them from performing the normal activity during the past 12 months or for a short and temporary period of 7 days. Compilation of the responses was aided by an illustrative body map to indicate the

major nine symptom sites - neck, shoulder, upper back, elbow, low back, wrist/hands, hip/thighs, knees, and ankles/ feet.

RESULT

The descriptive statistical analysis of data (N=947, Farmers), showed that the mean age was 47.83 ± 13.55 years. Figure 1 shows the musculoskeletal disorders in all 9 body regions mentioned in Self Developed Questionnaire and Figure 2 shows overall prevalence of WMSDs in farmers of Gujarat.




Figure 1: Prevalence of Work Related Musculoskeletal Disorders in farmers




Figure 2: Prevalence of Work Related Musculoskeletal Disorders in various regions of body in farmers

Out of the 947 study subjects, 80% have musculoskeletal disorders (MSDs) during a 12 month period as shown in Figure 1.

Figure 2 shows the distribution of musculoskeletal disorders in different parts of the body in male and female subjects according to Self developed questionnaire. The most common MSDs were found to be in the Lower back (61%) followed by Knee (33%), Shoulder (31%), Neck(29%), Upper back (28%), Elbow (15%) , Hip (13%), Wrist and hand (12%), and Ankle and Foot (11%).

DISCUSSION

Musculoskeletal disorders and the accompanying pain in farmers are multifactorial phenomena. Any part of the body is prone to develop such pain due to the work setup they are in and the nature of the work. Farmers in Gujarat work in various fields do a lot of manual labor. They also don't have knowledge of ergonomics which makes them susceptible to work related MSDs.

There are many types of research which document the prevalence of various musculoskeletal discomfort in an occupation like mine workers, stone cutters, sanitary workers, military personnel, aircrew workers, shoe factory workers, goldsmiths, etc. [13]

In the present study, farmers presented with pain in all the nine areas of the body identified in the Self Developed Questionnaire. But Low Back pain was the most common MSD that has been seen among the participants of the present study. Knee pain was the next common problem identified and these had been corroborated by multiple other studies carried out elsewhere i.e. Osborne et al, Murthy SR et al, Omran A et al. [15,27,28] The Neck and Shoulder pain were the other important MSDs affecting the farmers. The ankle was the region that was least affected among the workers in the study. However, these five areas were identified as most common musculoskeletal disorders areas- Lower Back(61%) followed by Knee (33%), Shoulder (31%), Neck(29%) and Upper back (28%).

Forward bending, twisting, exposure to vibrations while driving tractors coupled with heavy load carrying predispose them to continued physical stresses that affect the spinal disc making them prone to injury. Modern machinery like tractors, power tillers though may overtly seem to ease the work but actually put the workers to the additional risk of whole body vibrations. Walker-Bone and Palmer in their study on MSDs in agricultural workers concluded that tractor drivers or riders are particularly at risk of low backache for the concerned risk factors. [29, 30]

The present study also revealed substantial proportion of workers with knee involvement. The cultivators need to assume prolonged squatting positions in the fields that put excessive pressure on knee joints. Activities like squatting involve eccentric contraction of quadriceps muscle group. A study on agriculture health in University of Wisconsin USA reported that eccentric contraction leads to non-uniform lengthening of sarcomere or other ultra-structural abnormalities. [30, 31] These could possibly explain the development of knee pain in these workers.

Some ergonomic problems of farmers are twisting, bending, manual

material handling, awkward postures, and lifting, carrying heavy loads, hand tools, work-rest schedule and also lack of training of workers. [22-24] Such risk factors are associated with various musculoskeletal disorders. The risk of slipping, tripping, and fall on uneven fields is also associated with farming and these could also lead to the development of musculoskeletal discomfort in farmers. [25,26]

Long duration of involvement in farming activities was associated with higher prevalence of MSDs that had been corroborated by studies done by Osborne et al, Omran A et al., Xi. [4,28] Once pain in a particular body part developed, it was bound to remain over a long time and becomes chronic. These could have happened because of negligence or unavailability of proper health care facilities as well as lack of ergonomic education among farmers.

CONCLUSION

The prevalence of MSDs in farmers is very high. Low back pain is the most common type of MSD followed by Knee, Shoulder, Neck, Upper back, Elbow, Hip, Wrist and hand, and Ankle and Foot.

REFERENCES

1. Dr. Shamik PP, Dr. Kaushik M et al. Prevalence of Musculoskeletal Disorders and its Correlates among Agricultural Workers in Bhatar Block of Purba Bardhaman District, West Bengal.. *Journal of Dental and Medical Sciences*. January. 2019; 18(1):
2. Anamai T, Parvena M, Teerayut S. A Cross-sectional Study of Musculoskeletal Symptoms and Risk Factors in Cambodian Fruit Farm Workers in Eastern Region, Thailand. *Journal of Safety and Health at Work*. June 2018;9(2), 192-200.
3. <https://academic.oup.com/occmed/article-abstract/52/8/441/1404412> Musculoskeletal-disorders-in-farmers-and-farm workers by guest on 15 September 2017
4. Osborne A, Blake C, McNamara J, Meredith D, Phelan J, et al. Musculoskeletal disorders among Irish farmers. *Occup.Med (Lond)*. 2010; 60: 598-603.

5. Bihari V, Kesavachandran C, Pangtey BS, Srivastava AK, Mathur N. Musculoskeletal pain and its associated risk factors in residents of National Capital Region. *Indian J Occup Environ Med* 2011;15: 59-63.
6. Woolf AD, Pfleger B. Burden of major musculoskeletal conditions. *Bull. World Health Organ* 2003; 81: 646-656.
7. Allison TR, Symmons DP, Brammah T, Haynes P, Rogers A, et al. Musculoskeletal pain is more generalised among people from ethnic minorities than among white people in Greater Manchester. *Ann Rheum Dis* 2002; 61: 151-156.
8. Ghasemkhani M, Mahmudi E, Jabbari H. Musculoskeletal symptoms in workers. *Int J Occup Saf Ergon* 2008; 14: 455-462.
9. Dul J and Weerdmeester B. *Ergonomics for Beginners*. United States: CRC Press; 2001.
10. Kar SK, Dhara PC. An evaluation of musculoskeletal disorder and socioeconomic status of farmers in West Bengal, India. *Nepal Med Coll J*. 2007;9:245-49.
11. Bihari V, Kesavachandran C, Pangtey BS, Srivastava AK, Mathur N. Musculoskeletal pain and its associated risk factors in residents of National Capital Region. *Indian J Occup Environ Med*. 2011;15:59-63.
12. Allison TR, Symmons DP, Brammah T, Haynes P, Rogers A, et al. Musculoskeletal pain is more generalised among people from ethnic minorities than among white people in Greater Manchester. *Ann Rheum Dis*. 2002;61:151-56.
13. Birabi BN, Dienne PO, Ndukwu GU. Prevalence of low back pain among peasant farmers in a rural community in South Nigeria. *Rural Remote Health* 2012;12:1920.
14. Taechasubamorn P, Nopkesorn T, Pannarunothai S. Prevalence of low back pain among rice farmers in a rural community in Thailand. *J Med Assoc Thai*. 2011;94:616-21.
15. Osborne A., Blake C, Meredith D et al. Work-related musculoskeletal disorders among Irish farm operators. *Am. J. Ind. Med.*, 2013, 56: 235-242.
16. Safety and health in agriculture. Programme on Safety, Health and the Environment. Labour protection Department. International Labour Office. Geneva. http://www.ilo.org/wcmsp5/groups/public/-ed_protect/-protrav/
17. International Labour Organization. Sectoral Activities Programme. Code of practices on safety and health in agriculture. 2010. International Labour Office. Geneva http://www.ilo.org/wcmsp5/groups/public/-dgreports/-dcomm/-publ/documents/publication/wcms_159457.pdf. Last accessed on 19.01.2018
18. Rural, urban distribution of population. Census of India 2011. Registrar general and Census commissioner. Ministry of Home Affairs, New Delhi, India. http://censusindia.gov.in/2011-provresults/paper2/data_files/india/Rural_Urban_2011.pdf. Last accessed on 19.01.2018
19. Jha B. Employment, wages and productivity in Indian agriculture. Institute of Economic Growth, 2006, University of Delhi Enclave, Delhi, India.
20. Carol M. Black, Sickness absence and musculoskeletal disorders. *Rheumatology (Oxford)* 2012; 51 (2): 204-205.
21. Laxmaiah Manchikanti. Epidemiology of Low Back Pain. *Pain Physician, Association of Pain Management Anaesthesiologists* 2000; 3 (2):167- 192
22. Naeini HS, Karupiah K, Tamrin SB, Dalal K. Ergonomics in agriculture: An Approach in Prevention of Work-related Musculoskeletal Disorders (WMSDs). *Journal of Agriculture and Environmental Sciences*. 2014 June;3(2):33-51.
23. Parekh SK, Pathak NR. A comparative study between ergonomic advices versus ergonomic plus physiotherapy intervention in Low back pain among farmers. *Int J Physiother Res*. 2014;2(5):719-24.
24. Holmberg S, Stiernström EL, Thelin A, Svärdsudd K. Musculoskeletal symptoms among farmers and non-farmers: a population-based study. *Int J Occup Environ Health*. 2002 Oct-Dec;8(4):339-45.
25. Holmberg S, Thelin A, Stiernström EL, Svärdsudd K. Low back pain comorbidity among male farmers and rural referents: a population-based study. *Ann Agric Environ Med*. 2005;12:261-68.
26. Kolstrup CL. Work-related musculoskeletal discomfort of dairy farmers and employed workers. *J Occup Med Toxicol*. 2002;7:23.
27. Murthy SR, Nikhade N. Prevalence of musculoskeletal disorders in farmers of

- Ahmednagar district. *International J of Innovative Research in Medical Science* 2017;2(3):635-641
28. Omran A, Rezza G et al. Prevalence of musculoskeletal disorders among farmers in eastern Azerbaijan, Iran. *Indian J of Science and Technology*, 2015;8(28).
29. Walker-Bone K, Palmer KT. Musculoskeletal disorders in farmers and farm workers. *Occup. Med (London)*, 2002; 52(8): 441-450.
30. Hemalatha K, Bharanidharan G, Anusha T. Prevalence of musculoskeletal disorders among agricultural workers in rural area of Tamil Nadu: A cross-sectional study. *HECS International Journal of Community Health and Medical Research* 2017; 3(3):26-31
31. Sesto M. Chronic Musculoskeletal Disorders in Agriculture for partners in Agricultural Health. *Partners in Agricultural Health Module VIII*, University of Wisconsin, Madison, USA.

How to cite this article: Gadhavi B, Shukla Y. Prevalence of work related musculoskeletal disorders in farmers of Gujarat. *International Journal of Research and Review*. 2019; 6(11):231-236.
