

A Review on Development of Fast Dissolving Tablet and Approaches in Pharmaceutical Dosage of Atorvastatin

Dr. Rajesh Asija¹, Karishma Sharma², Seema Yadav Trimukhe³

^{1,2,3}Maharishi Arvind Institute of Pharmacy, Jaipur-302020, Rajasthan, India

Corresponding Author: Karishma Sharma

DOI: <https://doi.org/10.52403/ijrr.20220504>

ABSTRACT

Dosage form tablets and capsules in the current era still have gained considerable importance and acceptance by the formulation scientists, physicians, and the patients due to their numerous advantages which include easy self-administration and cost-effectiveness experienced by the patients. However, the drug delivery through the tablet dosage is no doubt, associated with potential drawbacks which include difficulty for the older and pediatric patients to swallow (dysphagia), low fraction of drug available for absorption in the bloodstream due to first-pass hepatic effect, and a slower therapeutic outcome in patients. Fast dissolving tablets have been emerged as a newer concept, in novel drug delivery systems that offer not only an alternative to conventional tablet dosage form, but also provide better therapeutic outcome, the ultimate solution, and a better option for the problem faced by the patients of such age group and diseases.

Keywords: Pharmaceutical dosage form, fast dissolving tablet, atorvastatin

INTRODUCTION

Pharmaceutical dosage forms and drug delivery systems are designed by formulation scientists as well as by various pharmaceutical manufacturing companies so that the required amount of the medicament can be delivered to the patients in a safer, precise, and effective manner. For this reason, several challenges and strategies are adopted and faced during the development

phase of such formulations to make them targeted, efficient, and stable throughout the shelf life period of the drug product.¹ Among all the drug delivery system presented in terms of oral solid dosage form tablets and capsules in the current era still have gained considerable importance and acceptance by the formulation scientists, physicians, and the patients due to their numerous advantages which include easy self-administration and cost-effectiveness experienced by the patients.² However the drug delivery through the tablet dosage is no doubt, associated with potential drawbacks which include difficulty for the older and pediatric patients to swallow (dysphagia), low fraction of drug available for absorption in the bloodstream due to first-pass hepatic effect, and a slower therapeutic outcome in patients.³

Fast Dissolving Tablets

Fast and rapid drug delivery to the site of action with minimum side effects has attracted pharmaceutical industries to formulate and designed a tablet dosage form that can dissolve, disintegrate in the mouth cavity, or disperses in water as quickly as possible. Thus readily absorbs into the bloodstream.⁴ According to a survey fifty percent of the population belonging to the class of geriatric and pediatric population find it difficult to swallow the tablet dosage form.⁵ So, for this purpose fast dissolving tablets have been emerged as a newer

concept, in novel drug delivery systems that offer not only an alternative to conventional tablet dosage form, but also provide better therapeutic outcome, the ultimate solution, and a better option for the problem faced by the patients of such age group and diseases.⁶

Approaches for Development

Following are the two approaches utilized:

Mouth dissolving tablets and Water dispersible tablets

Mouth Dissolving Tablets

This approach works by melting or disintegrating in the mouth cavity or by dissolving orally without the need of intake of water.⁷

Water Dissolving Tablets

This approach is designed and formulated in a way to disperse rapidly in water to form a homogenous dispersion.⁸

Reasons for Formulating Fast Dispersible Tablets

- Taste masking by the incorporation of various flavoring and sweetening agents.
- Enhanced patient compliance due to acceptability by the patients.
- The faster rate of disintegration in a matter of seconds.
- Ease of manufacturing with low cost.
- Offers the advantage of being stable due to changes in humidity and temperature.⁹
- Provides the benefit in targeting pediatric and geriatric populations to swallow both with and without the need of using water.¹⁰

Potential Candidates for Fast Dissolving Tablets

Following are the ideal characteristics that should be incorporated in the formulation development of fast dissolving tablets.

- The drug should exhibit better solubility in both water and saliva as compared to others.

- The fast dissolving tablet must cross, diffuse, and should be able to enter by means of partitioning into the epithelial lining of the upper part of GIT with (log P value be greater than one, or preferably greater than two).
- Must have a shorter biological half-life and frequent dosing.
- It should be able to permeate the tissue of oral mucosa.
- Molecular weight of the drug must be moderate to smaller in size.¹¹

Factors Influencing Formulation

Development of Fast Dissolving Tablets

- Physical and chemical properties of the active ingredient.
- Both the type and nature of active ingredients are incorporated.
- The method employed during manufacturing.¹²
- The physical and chemical properties of the excipients play an important role and therefore should be evaluated before formulation design and manufacturing.
- The design of formulation and the technique employed in manufacturing should be taken into consideration in order to avoid fluctuation in blood levels.

Salient Features and Developmental Challenges in The Development Of Fast Dissolving Tablets

Taste Masking

The unpleasant or bitter taste of the drug is a critical factor as the drug after disintegration into fine particles or granules in the oral mucosa is exposed to the taste buds. Hence, taste is one of the key parameters taken into account in development phase of fast dissolving drug delivery system. It is difficult to mask the bitter taste by the incorporation of sweetening agents alone readily available in the market. Therefore, in the recent years advancement in the technology has led to make improvement in such situations using combination of sweeteners with encapsulation and utilizing

the technique of coating and making granules.^{13, 14}

Moisture Sensitivity

Various fast dissolving tablets are moisture sensitive and are unstable to maintain the desired characteristic features of the drug under conditions of optimum humidity and controlled temperature. Therefore, several strategies are taken into consideration such as the addition of additives (water-soluble) to accelerate the dissolving ability and also enhance the taste. The addition of these additives imparts a serious damaging effect therefore; incorporation of suitable desiccants besides taking precautionary steps can prevent the deleterious effect of the hazards associated with the environment.^{15, 16}

Mouth Feel

Fast dissolving tablets must not break into granules in the mouth cavity. Therefore, particle size in the smaller range is preferred for formulation development. In addition, the property of mouthfeel can be achieved by the addition of cooling and flavoring agents that could result in enhanced patient compliance.¹⁷

Mechanical Strength And Disintegration Time

Since faster and quick disintegration activity with rapid onset of action has always been demanded by the patients. Several means in the past few decades were adopted to formulate it using different techniques like freeze-drying, wet mass compression, and molding that result in the formation of highly porous mass.^{18, 19} During the formulation development binder is added to provide mechanical strength and several water-soluble polymers like gelatin, dextran, and maltodextrin incorporated to maintain the shape. Secondly, agents like sucrose and mannitol facilitate the disintegration activity of the matrix supporting mediator.²⁰

Tablet Size

Patient compliance is also dependent on the size of the tablet which in turn is related to the patient comfort in intake of the dosage form. Research reported that the size of the tablet easier for the patient to swallow is 7-8 mm while the size preferred to handle is found to be larger than 8 mm. Therefore, both parameters of tablet size that encompass both easy swallowing and easy manufacturing are so far, quite problematic to accomplish.²¹

Methods Fast Dissolving Product Manufacturing Techniques

Direct Compression

Direct compression is the most preferable, demanding and simplest technique employed to manufacture a solid oral dosage form. Minimum number of steps involved, low cost of equipment utilized and commonly used excipients are the topmost features in selection of tablet manufacturing by direct compression technique.²² The technique involves a two step process in which the active (API) and excipients is mixed together and compacted using tableting machine.²³

Sublimation

Sublimation technique involves the formation of pores through the use of several volatile substances like thymol, camphor and menthol. The porous network of the matrix created results in quick water uptake which is due to the wicking action, hence faster rate of disintegration achieved.^{24, 25}

Moulding

It is another method to manufacture tablet in which either heat or solvent is used. The advantage of this method is better taste masking and quicker breaking up of tablet into fine particles or granules. However, these features are enhanced when porous mass of such tablets is produced using various physically modified components utilized in the procedure. The porous network created in the tablet is one of

the major steps in providing faster dissolution. However, several drawbacks associated with using this technique is poor mechanical strength that results in difficulty in handling. 26, 27

Freeze Drying

Freeze drying is also known as “Lyophilization”. In this process, the solvent (water) is removed from a solution or suspension form of a drug along with the carrier incorporated. Tablets manufacturing by freeze-drying produces tablets of high porosity and therefore, offer faster dissolution and disintegration. The medicament is released immediately as soon as it is kept in the oral cavity. The process is carried out at a temperature not high enough so as to reverse the effect produced by heat on the drug product, which will eventually improve the stability of the product. However, the major drawback of this technique is the high cost involve and also the product manufacture through freeze-drying is difficult to maintain the optimum feature of humidity and temperature.28

Melt Granulation

It is a process of manufacturing tablets in which powder mass is agglomerated using a binder that melts. The technique offers the advantage that no water or organic solvents is required. Since no drying step is involved, the process is less tedious, and therefore less energy than other granulation techniques are needed. It is a versatile technique to improve the dissolution profile of drugs.29

Super disintegrates Used In Fast Dissolving Tablets

Disintegrates play a pivotal role in facilitating the breaking up of a tablet into fine particles or granules when exposes to the medium (water). Hence, promote the water molecules to enter into the tablet and thus, mechanism of disintegration is achieved.30

The general mechanism of tablet disintegration is as follows.31

- Wetting down of tablet.

- The liquid to be dissolved enters into the crevices or pores.
- The added disintegrant swells due to absorption of water.
- Due to swelling, tablet disintegrates into powder particles or granules.

In order to produce robust tablets with desired properties of lesser disintegration and faster dissolution time particles of cohesive nature are required. The incorporated medicament must diffuse, dissolve and so a strong tablet resists dissolution. Tablet breaking agents added in the formulation to aid in the process of tablet disruption and fragmentation when exposes to water.32 Researchers has studied the effect, concentration and activity of disintegrants and super disintegrants on disintegration time. According to Rudinic and Schwartz, The concentration of starch suggested is 5% but this concentration can be increased by up to 15% if a drug has to been quickly disintegrated.33 In other study Davies studied the affect of hydrophobic lubricant on the activity of disintegrant added.34

Mechanism of Disintegrant Activity Water Wicking

Wicking through crospovidone is one of the effective mechanisms of disintegration and is defined as the ability of the disintegrant to penetrate water uptake into the tablet matrix resulting in the displacement of trapped air.35 However, porous tablets impart a significant effect in the disintegration pathway. The fluid enters into the crevices of the tablet matrix by means of capillary action and disrupts the interparticulate bond through which the tablet breaks.36

Swelling

Swelling is the mechanism of tablet deaggregation in which water uptake is facilitated in to the tablet matrix that causes disintegrants to swell and breaks.37 The porosity and packing fraction also imparts a significant impact on the disintegration

behavior therefore, powder with poor porosity exhibits poor disintegration time.³⁸

Repulsion

According to the theory based on the mechanism of particle repulsion tablet disintegration is also caused by particles that do not have the ability to swell. However, the tablet deaggregation mechanism depends on the forces of repulsion that exists between the powder particles. Hence researchers concluded that wicking is thought to be the primary mechanism of tablet breakup besides repulsion.³⁹

Particle Deformation

Disintegrants during the compression phase of tableting under deformation and particles return to their precompression shape upon wetting with water. Hence the mechanism of tablet deaggregation is achieved as number of deformed particle increases.⁴⁰

Effervescences

Tablet disintegration by effervescence involves the expulsion of carbon dioxide gas when a chemical reaction between citric and tartaric acid with bicarbonates and carbonates takes place. This favors the faster rate of dissolution with an added advantage of being good mouth feel.⁴¹

Enzymatic Reaction

The human body has several enzymes. These Enzymes help to break the bonds that exist between the primary powder particles. Hence, facilitates in the deaggregation mechanism of tablets.^{42, 43}

Atorvastatin

Atorvastatin (ATV), a HMG CoA reductase inhibitor, lipid lowering agent given in the dose ranging from 10-80 mg/day by oral route. After oral administration alone, ATV is rapidly absorbed; maximum plasma concentrations occur within 1 to 2 hours. Extent of absorption increases in proportion to atorvastatin dose. The absolute bioavailability of atorvastatin

(parent drug) is approximately 14% and the systemic availability of HMG-Co a reductase inhibitory activity is approximately 30%. The low systemic availability is attributed to pre-systemic clearance in gastrointestinal mucosa and/or hepatic first-pass metabolism.

Atorvastatin, a drug with low solubility and high permeability belongs to the Class II in the biopharmaceutics classification system (BCS), in which, dissolution process is the rate-limiting step for the absorption.^[44-45] Hence, it is important to drug features, such as the presence of polymorphism, stability, and compatibility of the pharmaceutical formulation, since any changes can directly influence its bioavailability.

CONCLUSION

The concept of a fast-dissolving drug delivery system emphasizes fulfilling the need of the patient in terms of providing faster onset of action. Because of such reasons patients found conventional drug delivery systems less easy, and less effective which result in decreased compliance issues related to swallowing. Hence, the concept of fast-dissolving drug delivery has been found as a newer approach that produces a significant impact in providing profitability and improving the life cycle of the drug product. It is, therefore, expected that new design future trends in such a drug delivery system will hopefully provide several different disciplines of current technology that would help to mitigate patients suffering from serious pain conditions.

Acknowledgement: None

Conflict of Interest: None

Source of Funding: None

REFERENCES

1. Chang RK, Guo X, Burnside BA, Couch RA. Fast-dissolving tablets. Pharm Tech Eur 2000; 12(6): 52–8.

2. R. Pahwa, M. Piplani, P. C. Sharma, D. Kaushik, S. Nanda. Orally disintegrating tablets friendly to pediatrics and geriatrics. *Archives App Sci* 2002; 2(2): 35–48.
3. Patil SL, Mahaparale PR, Shivnikar MA, Tiwari SS, Pawar KV, Sane PN. Fast dissolving oral films. An innovative drug delivery system. *Int J Res Rev Pharm App Sci* 2(3):482-96.
4. Prabhu SC, Parsekar SD, Shetty A, Monteiro SS, Azharuddin M, Shabaraya AR. A review on fast dissolving sublingual films for systemic drug delivery. *Int J Pharma Chem Sci* 2014; 3(2):501-11.
5. Mohanachandran PS, Krishna MPR, Fels S, Bini KB, Beenu B, Shilna KK. Formulation and evaluation of mouth dispersible tablets of amlodipine besylate. *Int J Appl Pharma* 2014; 2(3): 1-6.
6. Sanket K, Shiv KG. Fast dissolving tablets (FDTs) current status, new market opportunities, recent advances in manufacturing technologies and future prospects. *Int J Pharm Pharm Sci* 2014; 6(7):22-35.
7. Basu B, Bagadiya A, Makwana S, Vipul V, Batt D, Dharamsi A. Formulation and evaluation of fast dissolving tablets of cinnarizine using superdisintegrants blends and subliming material. *J Adv Pharm Technol Res* 2014; 2(4): 266-73.
8. Schiermeier S, Schmidt PC. Fast dispersible Ibuprofen tablets. *Eur J Pharm Sci* 2002; 15: 295-305.
9. Konar S, Mukhopadhyay A. Fast Dissolving drug delivery system. A novel approach. *Int J Pharm Biosci* 2014; 1(1): 1-10.
10. Mizumoto T, Masuda Y, Takeshi Y, Estuo Y and Katsuhide T. Formulation design of a novel fast disintegrating tablet. *Int J Pharm* 2005; 306 (12): 83–90.
11. Velmurugan S, Vinushitha S. Oral disintegrating tablets: An overview. *Int J Chem Pharm Sci* 2010; 1 (2): 1-12.
12. Amit KV, Anupam KS. A review on fast dissolving tablet as an efficient technique for oral drug delivery. *J Chem Pharm Sci* 2013; 6 (1): 29-34.
13. Sohi H, Sultana Y, Khar RK. Taste masking technologies in oral pharmaceuticals: Recent developments and approaches. *Drug Dev Ind Pharm* 2004; 30(5):429–48.
14. Ayenew Z, Puri V, Kumar L, Bansal AK. Trends in pharmaceutical taste masking technologies: A patent review. *Recent Pat Drug Deliv Formul* 2009; 3 (1): 26–39.
15. Chang RK, Guo X, Burnside BA, Couch RA. Fast-dissolving tablets. *Pharm Tech Eur* 2000; 12 (6): 52–8.
16. Katzner L, Jones B, Khattar J, Kosewick J. Blister package and packaged tablet. *US Patent* 2000; 6,155,423.
17. Bhandari S, Mittapalli RK, Gannu R, Rao YM. Orodispersible tablet. An overview. *Asian J Pharm* 2008; 2-10.
18. Jinichi F, Etsuo Y, Yasuo Y, Katsuhide T. Evaluation of rapidly disintegrating tablets containing glycine and carboxymethylcellulose. *Inter J Pharm* 2006; 310:101–9.
19. Segar H. Drug delivery products and the Zydis fast dissolving dosage. *J Pharm Pharm* 1998; 50 (4): 375–83.
20. Chandrasekhar R, Hassan Z, AlHusban F, Smith A, Mohammed A. The role of formulation excipients in the development of lyophilized fast-disintegrating tablets. *Eur J Pharm Biopharm* 2009; 72:119–29.
21. Sharma D, Kumar D, Singh M, Singh G, Rathore MS. Fast Disintegrating Tablets A new era in novel drug delivery system and new market opportunities. *J Drug Deliv Ther* 2012; 2(3): 74-86.
22. Tiwari N. A review on: Formulation and evaluation of fast dissolving tablet. *Int J Adv Res Pharm Bio Sci* 2013; 3(1): 60-9.
23. Harbir K. Processing Technologies for Pharmaceutical Tablets: A Review. *Int Res J Pharm* 2012; 3 (7): 20-3.

24. Shirsand SB, Suresh S, Kusumdevi V, Swamy PV. Formulation design and optimization of fast dissolving clonazepam tablets by sublimation method. *Indian J Pharm Sci* 2011; 73 (5): 491-6.
25. Mothilal M, Kumar AH, Krishna MC, Manasa V, Manimaran V, Damodharan N. Formulation and evaluation of modafinil fast dissolving tablets by sublimation technique. *J Chem Pharm Sci* 2013; 6 (3):147-54.
26. Dey P, Maiti S. Orodispersible tablets. A new trend in drug delivery. *J Nat Sci Biol Med* 2010; 1(1): 2-5.
27. Ghosh T, Ghosh A, Prasad D. A review on new generation orodispersible tablets and its future prospective. *Int J Pharm Pharm Sci* 2011; 3 (1): 1-7.
28. Fu Y, Yang S, Jeong SH, Kimura S, Park K. Orally fast disintegrating tablets. Developments, technologies, taste-masking and clinical studies. *Crit Rev Ther Drug Carrier Syst* 2004; 21 (6):433-76.
29. Dey P and Maiti S. Orodispersible tablets. A new trend in drug delivery. *J Nat Sci Biol Med* 2010; 1(1): 2-5.
30. Mangal M, Thakral S, Goswami M, Ghani P. Superdisintegrants. An updated review. *Int J Pharm Pharma Sci Res* 2012; 2(2):26-35.
31. Carstensen, J. T. 1993. Tablets. In: Carstensen, J.T, *Pharmaceutical Principles of Solid Dosage Forms*. Technomic publishing Co., Inc., Lancaster, Basel, pp. 86.
32. Armstrong, N.A. 2002. Tablet Manufacture In: Swarbrick, J., and Boylan. J.C. (Ed.), *Encyclopedia of Pharmaceutical Technology* 2002; 2nd Edn., Vol, 03. Marcel Dekker, Inc., New York, Basel, pp.2721.
33. Rudinic, E.M. and Schwartz, J.B., 2006. Oral Solid Dosage Forms. In: Gennaro, A. R. (Ed.), *Remington: The Science and practice of Pharmacy*, 21st Edn., Lippincott Williams and Wilkins, A wolters kluwer company, Philadelphia, Baltimore, New York, London, Bueenos Aires, pp.893,901.
34. Davies, P., 2001. Oral Solid Dosage Forms. In: Gibson, M. (Ed.), *Pharmaceutical Preformulation and Formulation*. Taylor and Francis, Boca Raton, London, New York, Singapore, pp. 419-420.
35. Augsburger, L.L., Hahm, H. A., Brzecko, A. W. and Shah, S. 2002. Super Disintegrants: Characterization and Function In: Swarbrick, J., and Booylan, J.C. (Ed.), *Encyclopedia of Pharmaceutical technology*; 2nd Edn., vol. 03. Marcel Dekker, Inc., New York, Basel. Pp. 2623-2635.
36. Mangal M, Thakral S, Goswami M, Ghai P. Superdisintegrants. An updated review. *Int J Pharm Pharm Sci Res* 2012; 2(2): 26-35.
37. Khairnar DA, Anantwar SP, Chaudhari CS, Shelke PA. Superdisintegrants. An emerging paradigm in orodispersible tablets. *Int J Biopharm* 2014; 5:119-28.
38. Bala R, Khanna S, Pawar P. Polymers in fast disintegrating tablets. a review. *Asian J Pharm Clin Res* 2012; 5 (2): 8-14.
39. Dass S, Mazumedar SA. An updated Precise Review on Super disintegrants. *Indo Am J Pharm Res* 2013; 3(6): 5342-55.
40. Roy D, Bhowmik D, Kumar KS. A comprehensive review on superdisintegrants used in Orodispersible tablets. *Indian J Res Pharm Biotechnol* 2014; 2(4): 1297-303.
41. Roy A, Orodispersible Tablets: A review. *Asian J Pharm Clin Res* 2016; 9(1): 19-26.
42. Kavitha K, Hui BJ, Subramaniam K, Santhi K, Dhanaraj SA, Rupeshkumar M. Role of superdisintegrants in the design of fast dissolving drug delivery systems. A review. *Int J Adv Pharm Res* 2013; 4(11): 2475-83.
43. Vimal VV, Aarathi TS, Soumya BJ. Superdisintegrants in fast disintegrating drug delivery systems. A brief review. *Int J Pharm* 2013; 3(2): 380-385.

44. Rasheed A, Aishwarya K, Basha N, Reddy B.S, Swetha A. Dexibuprofen-dextran macromolecular prodrugs: Synthesis, characterization and pharmacological evaluation. *Der Pharma Chemica* 2009; (2):124–32.
45. Drug Bank 2017. Drug created on October 21, 2015 09:53 / Updated on August 02, 2017 17:33.

<https://www.drugbank.ca/drugs/DB01050>.

How to cite this article: Rajesh Asija, Karishma Sharma, Seema Yadav Trimukhe. A review on development of fast dissolving tablet and approaches in pharmaceutical dosage of atorvastatin. *International Journal of Research and Review*. 2022; 9(5): 15-22.
DOI: <https://doi.org/10.52403/ijrr.20220504>
